

INTERPRETING KEY TRIALS



EDUCATIONAL OBJECTIVE: Readers will engage their patients in a shared decision-making process regarding intensive lowering of blood pressure, with discussion of the benefits and risks

GEORGE THOMAS, MD, FACP, FASN

Director, Center for Blood Pressure Disorders, Department of Nephrology and Hypertension, Glickman Urological and Kidney Institute, Cleveland Clinic; Assistant Professor, Cleveland Clinic Lerner College of Medicine of Case Western Reserve University, Cleveland, OH; Cleveland Clinic Site Principal Investigator, Systolic Blood Pressure Intervention Trial (SPRINT)

JOSEPH V. NALLY, MD

Director, Center for Chronic Kidney Disease, Department of Nephrology and Hypertension, Glickman Urological and Kidney Institute, Cleveland Clinic; Clinical Professor, Cleveland Clinic Lerner College of Medicine of Case Western Reserve University, Cleveland, OH

MARC A. POHL, MD

Ray W. Gifford Chair in Hypertension, Department of Nephrology and Hypertension, Glickman Urological and Kidney Institute, Cleveland Clinic; Chair, Clinical Management Committee, Irbesartan Diabetic Nephropathy Trial

Interpreting SPRINT: How low should you go?

ABSTRACT

The Systolic Blood Pressure Intervention Trial (SPRINT) found evidence of cardiovascular benefit with intensive lowering of systolic blood pressure (goal < 120 mm Hg) compared with the currently recommended goal (< 140 mm Hg) in older patients with cardiovascular risk but without diabetes or stroke. This article reviews the trial design and protocol, summarizes the results, and briefly discusses the implications of these results.

KEY POINTS

SPRINT is the first large prospective randomized trial to show evidence of cardiovascular and mortality benefit for intensive lowering of systolic blood pressure (goal < 120 mm Hg) in older patients at cardiovascular risk, but without a history of diabetes mellitus or stroke. A similar trial in patients with type 2 diabetes mellitus did not show significant benefit of intensive treatment.

Intensive treatment was associated with more adverse events, including hypotension, syncope, electrolyte abnormalities, and acute kidney injury.

It is unclear if these results can be extrapolated to patients with a history of diabetes or stroke, younger patients, or those with low cardiovascular risk.

Healthcare providers should engage patients in a shared decision-making process, with discussion of the benefits and risks associated with intensive lowering of blood pressure.

IN TREATING HYPERTENSION, lower systolic pressure is better than higher—with caveats. This is the message of the Systolic Blood Pressure Intervention Trial (SPRINT),¹ a large, federally funded study that was halted early when patients at high cardiovascular risk who were randomized to a goal systolic pressure of less than 120 mm Hg were found to have better outcomes, including lower rates of heart failure, death from cardiovascular causes, and death from any cause, than patients randomized to a goal of less than 140 mm Hg.

See related editorial, page 196

The caveats: the benefit came at a price of more adverse events. Also, the trial excluded patients who had diabetes mellitus or previous strokes, so it is uncertain if these subgroups would also benefit from intensive lowering of systolic pressure—and in earlier trials they did not.

This article reviews the trial design and protocol, summarizes the results, and briefly discusses the implications of these results.

■ BEFORE SPRINT

Hypertension is very common in adults in the United States, and is a risk factor for heart disease, stroke, heart failure, and kidney disease. The estimated prevalence of hypertension in the 2011–2014 National Health and Nutrition Examination Survey (NHANES) was 29%, and the prevalence increases with age (7.3% in those ages 18 to 39, 32.2% in those ages 40 to 59, and 64.9% in those ages 60 and older).² Isolated systolic hypertension (ie, systolic blood pressure > 140 mm Hg with diastolic pressure < 90 mm Hg) is the most common form of hypertension after age 50.³

SPRINT was stopped early due to better outcomes in the intensive treatment group

Clinical trials have provided substantial evidence that treating hypertension reduces the incidence of stroke, myocardial infarction, and heart failure.^{4,5} Although observational studies show a progressive and linear rise in cardiovascular risk as systolic blood pressure rises above 115 mm Hg,⁶ clinical trials in the general population have not documented benefits of lowering systolic pressure to this level.⁷⁻¹¹ However, clinical trials that directly evaluated two different blood pressure goals in the general population showed benefit with achieving systolic blood pressure less than 150 mm Hg,^{7,9} with limited data on lower blood pressure targets.¹⁰⁻¹²

No benefit found in intensive systolic lowering in diabetes or after stroke

The Action to Control Cardiovascular Risk in Diabetes-Blood Pressure (ACCORD BP) trial¹³ in patients with type 2 diabetes found no benefit in lowering systolic pressure to less than 120 mm Hg compared with less than 140 mm Hg in terms of the trial's primary composite cardiovascular outcome (ie, nonfatal myocardial infarction, nonfatal stroke, or death from cardiovascular causes). However, the intensively treated group in this trial did enjoy a benefit in terms of fewer stroke events.

The Secondary Prevention of Small Subcortical Strokes (SPS3) trial¹⁴ in patients with stroke found no significant benefit in lowering systolic pressure to less than 130 mm Hg compared with less than 150 mm Hg for overall risk of another stroke, but a significant benefit was noted in reduced risk of intracerebral hemorrhage.

Current guidelines, based on available evidence, advocate treatment to a systolic goal of less than 140 mm Hg in most patients, and recommend relaxing this goal to less than 150 mm Hg in the elderly.^{15,16}

Given the uncertainty surrounding optimal systolic targets, SPRINT was designed to test the hypothesis that a goal of less than 120 mm Hg would reduce the risk of cardiovascular events more than the generally accepted systolic goal of less than 140 mm Hg.¹⁷ Patients with diabetes and stroke were excluded because a similar hypothesis was tested in the ACCORD BP and SPS3 trials, which included patients with these conditions.

SPRINT DESIGN

SPRINT was a randomized, controlled, open-label trial sponsored by the National Institutes of Health and conducted at 102 US sites.

Inclusion criteria. Participants had to be at least 50 years old, with systolic pressure of 130 to 180 mm Hg, and had to have at least one cardiovascular risk factor, eg:

- Clinical or subclinical cardiovascular disease (other than stroke)
- Chronic kidney disease, defined as estimated glomerular filtration rate (eGFR), calculated by the Modification of Diet in Renal Disease (MDRD) study equation, of 20 to less than 60 mL/min/1.73 m²
- Framingham risk score of 15% or more
- Age 75 or older.

Major exclusion criteria included:

- Diabetes
- Stroke
- Polycystic kidney disease
- Chronic kidney disease with an eGFR less than 20 mL/min/1.73 m²
- Proteinuria (excretion > 1 g/day).

Intensive vs standard treatment

Participants were randomized to receive intensive treatment (systolic goal < 120 mm Hg) or standard treatment (systolic goal < 140 mm Hg). Baseline antihypertensive medications were adjusted to achieve blood pressure goals based on randomization assignment.

Doses of medications were adjusted on the basis of an average of three seated office blood pressure measurements after a 5-minute period of rest, taken with an automated monitor (Omron Healthcare Model 907); the same monitor was used and the same protocol was followed at all participating sites. Blood pressure was also measured after standing for 1 minute to assess orthostatic change.

Lifestyle modifications were encouraged in both groups. There was no restriction on using any antihypertensive medication, and this was at the discretion of individual investigators. Thiazide-type diuretics were encouraged as first-line agents (with chlorthalidone encouraged as the primary thiazide-type diuretic).

Outcomes measured

The **primary outcome** was a composite of myocardial infarction, acute coronary syn-

drome not resulting in myocardial infarction, stroke, acute decompensated heart failure, and cardiovascular mortality.

Secondary outcomes included individual components of the primary composite outcome, all-cause mortality, and the composite of primary outcome and all-cause mortality.

Renal outcomes were assessed as:

- Incident albuminuria (doubling of the urinary albumin-to-creatinine ratio from less than 10 mg/g to more than 10 mg/g)
- Composite of a 50% decrease in eGFR or development of end-stage renal disease requiring long-term dialysis or kidney transplantation (in those with baseline chronic kidney disease)
- A 30% decrease in eGFR (in those without chronic kidney disease).^{1,17}

SPRINT also recruited participants to two nested substudies: SPRINT MIND and SPRINT MIND MRI, to study differences in cognitive outcomes and small-vessel ischemic disease between intensive treatment and standard treatment.

■ **STUDY RESULTS**

Older patients at risk, but without diabetes

Of 14,692 participants screened, 9,361 were enrolled in the study between 2010 and 2013. Baseline characteristics were comparable in both groups.

Demographics. The mean age of the participants was 67.9, and about 28% were 75 or older. About 36% were women, 58% white, 30% black, and 11% Hispanic.

Cardiovascular risk. The mean Framingham risk score was 20% (ie, they had a 20% risk of having a cardiovascular event within 10 years), and 61% of the participants had a risk score of at least 15%. Twenty percent already had cardiovascular disease.

Blood pressure. The average baseline blood pressure was 139.7/78.2 mm Hg. One-third of the participants had baseline systolic pressures of 132 mm Hg or less, another third had pressures in the range of 132 to 145, and the rest had 145 mm Hg or higher.

Renal function. The mean serum creatinine level was about 1.1 mg/dL. The mean eGFR was about 71 mL/min/1.73 m² as calculated by the MDRD equation, and about 28%

TABLE 1

Antihypertensive medications used in SPRINT

Medication class	Intensive therapy (%)	Standard therapy (%)
Angiotensin-converting enzyme inhibitors or angiotensin II receptor blockers	76.7	55.2
Thiazide-type diuretics	54.9	33.3
Dihydropyridine calcium channel blockers	52.8	31.3
Beta-blockers	41.1	30.8
Aldosterone antagonists	8.7	4.0
Other potassium-sparing diuretics	3.1	2.5
Nondihydropyridine calcium channel blockers	4.7	4.3
Direct vasodilators	7.3	2.4

Information from SPRINT Research Group; Wright JT Jr, Williamson JD, Whelton PK, et al. A randomized trial of intensive versus standard blood-pressure control. *N Engl J Med* 2015; 373:2103–2116.

had eGFRs less than 60. The mean ratio of urinary albumin to creatinine was 44.1 mg/g in the intensive treatment group and 41.1 in the standard treatment group.

Other. The mean total cholesterol level was 190 mg/dL, fasting plasma glucose 99 mg/dL, and body mass index nearly 30 kg/m².

Blood pressure during treatment

Throughout the 3.26 years of follow-up, the average difference in systolic pressure between the two groups was 13.1 mm Hg, with a mean systolic pressure of 121.5 mm Hg in the intensive treatment group and 134.6 mm Hg in the standard treatment group. The mean diastolic blood pressure was 68.7 mm Hg in the intensive treatment group and 76.3 mm Hg in the standard treatment group.

People in the intensive treatment group were taking a mean of 2.8 antihypertensive medications, and those in the standard treatment group were taking 1.8. Patients in the intensive group required greater use of all classes of medications to achieve goal systolic pressure (Table 1).

Study halted early due to efficacy

Although the study was planned to run for an average follow-up of 5 years, the National Heart, Lung, and Blood Institute terminated it early at a median of 3.26 years in view of low-

Intensive treatment required, on average, one more medication than standard treatment

TABLE 2

SPRINT results at a glance

Outcome	Percent per year		Hazard ratio
	Intensive therapy	Standard therapy	
Primary outcome^a	1.65	2.19	0.75 ^b
Secondary outcomes			
Myocardial infarction	0.65	0.78	0.83
Other acute coronary syndromes	0.27	0.27	1.00
Stroke	0.41	0.47	0.89
Heart failure	0.41	0.67	0.62 ^b
Cardiovascular mortality	0.25	0.43	0.57 ^b
Other secondary outcomes			
All-cause mortality	1.03	1.40	0.73 ^b
In patients with chronic kidney disease—decrease in eGFR of ≥ 50% or end-stage renal disease	0.33	0.36	0.89
In patients without chronic kidney disease—decrease in eGFR of ≥ 30% to < 60 mL/min/1.73 m ²	1.21	0.35	3.49 ^b
Adverse events	Percent of patients		Hazard ratio
	Intensive therapy	Standard therapy	
Hypotension	3.4	2.0	1.70 ^b
Syncope	3.5	2.4	1.44 ^b
Hyponatremia	3.8	2.1	1.76 ^b
Hypokalemia	2.4	1.6	1.50 ^b
Injurious fall	7.1	7.1	1.00
Orthostatic hypotension without dizziness	16.6	18.3	0.88 ^b
Orthostatic hypotension with dizziness	1.3	1.5	0.85
Acute kidney injury	4.4	2.6	1.71 ^b

^aThe composite of myocardial infarction, acute coronary syndrome, stroke, heart failure, or death from cardiovascular causes.

^b*P* < .05.

eGFR = estimated glomerular filtration rate, according to the Modification of Diet in Renal Disease study equation.

Information from SPRINT Research Group; Wright JT Jr, Williamson JD, Whelton PK, et al. A randomized trial of intensive versus standard blood-pressure control. *N Engl J Med* 2015; 373:2103–2116.

er rates of the primary outcome and of heart failure and death in the intensive treatment group (Table 2).

The effects on the primary outcome and mortality were consistent across the prespecified subgroups of age (< 75 vs ≥ 75), sex (female vs male), race (black vs nonblack), cardiovascular disease (presence or absence at baseline), prior chronic kidney disease (presence or absence at baseline), and across blood pressure tertiles (≤ 132 mm Hg, > 132 to < 145 mm Hg, ≥ 145 mm Hg).

Follow-up for assessment of cognitive outcomes (SPRINT MIND) and small-vessel ischemic disease (SPRINT MIND MRI) is ongoing.

WHAT DOES THIS MEAN?

SPRINT is the first large, adequately powered, randomized trial to demonstrate cardiovascular and mortality benefit from lowering the systolic blood pressure (goal < 120 mm Hg) in older patients at cardiovascular risk but without a history of diabetes mellitus or stroke.¹

Most SPRINT patients had reasonably controlled blood pressure at baseline (the mean systolic pressure was 139.7 mm Hg, and two-thirds of participants had systolic pressure < 145 mm Hg). Of note, however, this trial excluded patients with systolic pressure higher than 180 mm Hg. There was excellent separation of systolic pressure between the two groups beginning at 1 year, which was consistent through the course of the trial.

The cardiovascular benefit in the intensive treatment group was predominantly driven by lower rates of heart failure (a 38% reduction in the intensive treatment group, *P* = .0002) and cardiovascular mortality (a 43% reduction in the intensive treatment group, *P* = .005), while there was no significant difference between the two groups in myocardial infarction or stroke. The beneficial effect on heart failure events is consistent with results from other trials including the Systolic Hypertension in the Elderly Program,⁷ Systolic Hypertension in Europe,⁸ and Hypertension in the Very Elderly Trial,⁹ all of which showed greatest risk reduction for heart failure events with systolic pressure-lowering (although to higher systolic levels than SPRINT).⁷⁻⁹ It is unclear why there was no beneficial effect on stroke events. The reduction in all-cause mortality in the intensive treatment group in SPRINT

was greater than the reduction in cardiovascular deaths, which is also unexplained.

Although the study was terminated early due to efficacy (which introduces the possible bias that the estimated effect size will be too high), the number of primary end points reached was large (562 in the two groups combined), providing reassurance that the findings are valid. There was no blinding in the study (both participants and study investigators were aware of treatment assignment and study medications), but there was a structured assessment of outcomes and adverse events, with adjudication done by blinded reviewers.

SPRINT used an automated device for blood pressure measurement, which is known to reduce the “white coat” effect and correlates tightly with average daytime blood pressure done by ambulatory blood pressure monitoring.¹⁸ However, in clinical practice automated devices may not be available and a strict protocol for correct measurement may not be followed, with the possible result that blood pressure may be overestimated and overtreated.

What about diastolic pressure?

The trial, by design, focused on lowering systolic pressure (given the greater prevalence of isolated systolic hypertension with age), and the implications of lowering diastolic pressure are unclear. The issue of a J-shaped relationship between diastolic pressure and cardiovascular risk is debated in the literature: patients with a diastolic pressure of 60 to 65 mm Hg, especially those with existing coronary artery disease, may not tolerate aggressive blood pressure-lowering.^{19,20} Further analysis of this association (if any) from SPRINT will be helpful.

What about patients with diabetes?

Patients were excluded from SPRINT if they were under age 50, were at low cardiovascular risk, or had diabetes, raising the question of whether the results apply to these groups as well.

The question is particularly relevant in diabetes, as the ACCORD BP study, which used the same blood pressure targets as SPRINT, did not show a significant difference in the primary cardiovascular outcome between the intensive and standard treatments in patients with diabetes (Table 3).¹³ In ACCORD BP,

the rate of the primary outcome was 12% lower in the intensive treatment group than in the standard treatment group, but the 95% confidence interval was –27% to +6%, so the finding was not statistically significant. However, the wide confidence interval does not exclude the possibility of a benefit that was comparable to that observed in SPRINT.

It has been speculated that ACCORD BP was underpowered to detect significant differences in the primary outcome.²¹ An analysis combining data from both trials indicated that effects on individual outcomes were generally consistent in both trials (with no significant heterogeneity noted).²² Also, the primary composite outcome in ACCORD did not include heart failure, which is particularly sensitive to blood pressure reduction.

Additionally, ACCORD BP had a 2 × 2 factorial design involving a simultaneous comparison of intensive vs standard glycemic control, which may have influenced the effects due to blood pressure. Indeed, a post hoc analysis showed that there was a significant 26% lower risk of the primary outcome in ACCORD BP patients who received intensive systolic pressure control plus standard glycemic control than in those receiving standard systolic control plus standard glycemic control.²³

Are more adverse events an acceptable trade-off?

Adverse events, including acute kidney injury, were more frequent in the intensive therapy group in SPRINT.

Acute kidney injury was coded as an adverse event on the basis of this diagnosis being included in the hospital discharge summary (as a primary or main secondary diagnosis) and if considered by the safety officer to be one of the top three reasons for admission or continued hospitalization. Further analysis of renal events should be forthcoming.

People in the intensive treatment group, on average, needed one more medication than those in the standard treatment group. Some of the adverse events may be related to the anti-hypertensive medications taken (eg, electrolyte abnormalities such as hyponatremia and hypokalemia due to diuretic use), and others may be related to blood pressure-lowering (eg, acute

Only about half of hypertensive adults have their blood pressure under control, ie, < 140/90

TABLE 3

Comparing the ACCORD BP trial and SPRINT

	ACCORD BP¹³	SPRINT¹
Trial design	Randomized controlled trial, 2 × 2 factorial design—intensive vs standard antihyperglycemic treatment and intensive vs standard antihypertensive treatment	Randomized controlled trial, intensive vs standard antihypertensive treatment
Number of participants	4,733	9,361
Main inclusion criteria	Type 2 diabetes Systolic pressure 130–180 mm Hg Age 40 and older (upper age limit 79)	Systolic pressure 130–180 mm Hg Age 50 and older
Main exclusion criteria	Serum creatinine > 1.5 mg/dL	Stroke, diabetes mellitus
Follow-up	Mean 4.7 years	Median 3.26 years
Mean age	62.2	67.9
Female	47.7%	35.6%
Black	24.1%	29.9%
Baseline cardiovascular disease	33.7%	20.1%
Mean estimated glomerular filtration rate	91.6 mL/min/1.73 m ²	71.7 mL/min/1.73 m ²
Mean achieved systolic blood pressure	119.3 vs 133.5 mm Hg	121.5 vs 134.6 mm Hg
Mean achieved diastolic pressure	64.4 vs 70.5 mm Hg	68.7 vs 76.3 mm Hg
Mean number of medications	3.4 vs 2.1	2.8 vs 1.8
Diuretic of choice	Hydrochlorothiazide	Chlorthalidone
Primary outcome definition	Composite of nonfatal myocardial infarction, nonfatal stroke, and cardiovascular mortality	Composite of myocardial infarction, other acute coronary syndromes, stroke, heart failure, and cardiovascular mortality
Primary outcome results	1.87 vs 2.09%/year	1.65 vs 2.19%/year ^a
Individual outcome results		
Myocardial infarction	1.13 vs 1.28%/year	0.65 vs 0.78%/year
Stroke	0.32 vs 0.53%/year ^a	0.41 vs 0.47%/year
Cardiovascular mortality	0.52 vs 0.49%/year	0.25 vs 0.43%/year ^a
Heart failure	0.73 vs 0.78%/year	0.41 vs 0.67%/year ^a
All-cause mortality	1.28 vs 1.19%/year	1.03 vs 1.40%/year ^a
Adverse events		
Hypotension	0.7 vs 0.04% ^a	3.4 vs 2.0% ^a
Syncope	0.5 vs 0.21% ^a	3.5 vs 2.4% ^a
Hypokalemia	2.1 vs 1.1% ^a	2.4 vs 1.6% ^a

^a*P* < .05.

Information from ACCORD Study Group; Cushman WC, Evans GW, Byington RP, et al. Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med* 2010; 362:1575–1585 and SPRINT Research Group; Wright JT Jr, Williamson JD, Whelton PK, et al. A randomized trial of intensive versus standard blood-pressure control. *N Engl J Med* 2015; 373:2103–2116.

kidney injury due to renal hypoperfusion).

At this point, the long-term effects of these adverse events, especially on kidney function, are not known. Patients enrolled in clinical trials tend to be healthier than patients seen in clinical practice; thus, the rate of adverse events reported in the trial may be lower than one would see in the real world.

Does lower systolic pressure protect or harm the kidneys?

SPRINT included patients with stage 3 and 4 chronic kidney disease (ie, with eGFR 20–50 mL/min/1.73 m²), but it was designed to assess cardiovascular outcomes, not the progression of chronic kidney disease. The trial excluded patients with diabetic nephropathy or high degrees of proteinuria.

Earlier randomized trials that focused on chronic kidney disease progression, including the MDRD²⁴ and the African American Study of Kidney Disease and Hypertension,²⁵ did not show benefit with more aggressive blood pressure-lowering (except in patients with higher degrees of proteinuria), and these trials were not powered to assess effects on cardiovascular outcomes.^{24,25}

The Irbesartan Diabetic Nephropathy Trial,^{26,27} which was done in patients with overt diabetic nephropathy, showed that a progressively lower achieved systolic pressure down to 120 mm Hg predicted lower rates of heart failure, cardiovascular mortality, and renal events (although the trial target was ≤ 130/85 mm Hg and few participants achieved systolic pressure lower than 120 mm Hg).

■ IMPLICATIONS FOR MANAGEMENT

The recent estimates of hypertension prevalence and control from NHANES show that only about 53% of hypertensive adults have their blood pressure under control (defined as systolic pressure < 140 mm Hg and diastolic pressure < 90 mm Hg).² Analysis of the NHANES 2007–2012 data showed that 16.7% or 8.2 million US adults with treated hypertension meet the eligibility criteria for SPRINT.²⁸

Although the SPRINT results support the notion that “lower is better,” the risks and benefits of intensive control will need to be balanced in individual patients. **Table 4** shows

TABLE 4

Benefit vs harm of intensive and standard systolic pressure control in SPRINT

Outcome	Absolute risk reduction	Number needed to treat
Primary outcome	–1.6%	61
Heart failure	–0.8%	125
Cardiovascular mortality	–0.6%	167
All-cause mortality	–1.2%	83
	Absolute risk increase	Number needed to harm
Hypotension	+1.0%	100
Syncope	+0.6%	167
Electrolyte abnormalities	+0.8%	125
Acute kidney injury	+1.6%	62

Information from SPRINT Research Group; Wright JT Jr, Williamson JD, Whelton PK, et al. A randomized trial of intensive versus standard blood-pressure control. *N Engl J Med* 2015; 373:2103–2116.

the number needed to treat and number needed to harm in the trial.

More aggressive management of hypertension is challenging. The median systolic pressure achieved in the intensive group in SPRINT was just over 120 mm Hg, which implies that at least half of the participants in the intensive group did not achieve the goal of less than 120 mm Hg. While it may be reasonable to aim for systolic pressure of less than 120 or 125 mm Hg in patients who fit the SPRINT criteria *and* can tolerate intensive blood pressure lowering, it would be prudent to aim for a more conservative goal in elderly patients who are frail and at risk for falls, considering the higher incidence of specified adverse events in the intensive group.

Results of cognitive outcomes, as well as data related to quality of life, are still awaited. Long-term renal outcomes are also unclear.

As noted above, the question of generalizability of SPRINT results to patients with diabetes is open to debate. In our opinion, with currently available evidence, it is difficult to conclusively answer the question of whether a lower systolic target provides cardiovascular benefit in diabetes. It is also unclear whether similar beneficial results would be seen with

Lower systolic pressure is better if adverse events can be monitored, avoided, or managed

intensive treatment in a population at low cardiovascular risk. The American Heart Association and the American College of Cardiology are in the process of formulating new hypertension guidelines, and evidence from SPRINT will inform any new recommendations.

As more medications will likely be needed for intensive systolic blood pressure control, side effects and tolerability of medications with polypharmacy and potential nonadherence with increasing complexity of medication regimens should be kept in mind. Lifestyle modifications will need to be emphasized and reinforced, with greater use of combina-

tion antihypertensive therapy.

The data from SPRINT indicate that lower systolic pressure is better, as long as untoward clinical events can be monitored and avoided or easily managed. Careful monitoring will likely entail more frequent clinic visits and more frequent assessment of renal function and electrolyte levels (participants in the intensive group in the trial were seen every month until goal was achieved). A team approach that includes pharmacists and nurse practitioners, along with optimal use of best practice algorithms and remote monitoring technology, will need to be implemented for efficient and effective care. ■

REFERENCES

1. **SPRINT Research Group; Wright JT Jr, Williamson JD, Whelton PK, et al.** A randomized trial of intensive versus standard blood-pressure control. *N Engl J Med* 2015; 373:2103–2116.
2. **Yoon SS, Fryar CD, Carroll MD.** Hypertension prevalence and control among adults: United States, 2011–2014. NCHS data brief, no. 220. Hyattsville, MD: National Center for Health Statistics. 2015
3. **Franklin SS, Jacobs MJ, Wong ND, L'Italien GJ, Lapuerta P.** Prevalence of isolated systolic hypertension among middle-aged and elderly US hypertensives: analysis based on National Health and Nutrition Examination Survey (NHANES) III. *Hypertension* 2001; 37:869–874.
4. **Neal B, MacMahon S, Chapman N; Blood Pressure Lowering Treatment Trialists' Collaboration.** Effects of ACE inhibitors, calcium antagonists, and other blood-pressure-lowering drugs: results of prospectively designed overviews of randomised trials. *Blood Pressure Lowering Treatment Trialists' Collaboration.* *Lancet* 2000; 356:1955–1964.
5. **Psaty BM, Smith NL, Siscovick DS, et al.** Health outcomes associated with antihypertensive therapies used as first-line agents. A systematic review and meta-analysis. *JAMA* 1997; 277:739–745.
6. **Lewington S, Clarke R, Qizilbash N, Peto R, Collins R; Prospective Studies Collaboration.** Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 2002; 360:1903–1913.
7. **SHEP Cooperative Research Group.** Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension: final results of the Systolic Hypertension in the Elderly Program (SHEP). *JAMA* 1991; 265:3255–3264.
8. **Staessen JA, Fagard R, Thijs L, et al.** Randomised double-blind comparison of placebo and active treatment for older patients with isolated systolic hypertension. The Systolic Hypertension in Europe (Syst-Eur) Trial Investigators. *Lancet* 1997; 350:757–764.
9. **Beckett NS, Peters R, Fletcher AE, et al; HYVET Study Group.** Treatment of hypertension in patients 80 years of age or older. *N Engl J Med* 2008; 358:1887–1898.
10. **JATOS Study Group.** Principal results of the Japanese trial to assess optimal systolic blood pressure in elderly hypertensive patients (JATOS). *Hypertens Res* 2008; 31:2115–2127.
11. **Ogihara T, Saruta T, Rakugi H, et al; Valsartan in Elderly Isolated Systolic Hypertension Study Group.** Target blood pressure for treatment of isolated systolic hypertension in the elderly: Valsartan in Elderly Isolated Systolic Hypertension study. *Hypertension* 2010; 56:196–202.
12. **Liu L, Zhang Y, Liu G, Li W, Zhang X, Zanchetti A; FEVER Study Group.** The Felodipine Event Reduction (FEVER) Study: a randomized long-term placebo-controlled trial in Chinese hypertensive patients. *J Hypertens* 2005; 23:2157–2172.
13. **ACCORD Study Group; Cushman WC, Evans GW, Byington RP, et al.** Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med* 2010; 362:1575–1585.
14. **SPS3 Study Group; Benavente OR, Coffey CS, Conwit R, et al.** Blood-pressure targets in patients with recent lacunar stroke: the SPS3 randomised trial. *Lancet* 2013; 382:507–515.
15. **James PA, Oparil S, Carter BL, et al.** 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA* 2014; 311:507–520. Erratum in: *JAMA*. 2014; 311:1809.
16. **Weber MA, Schiffrin EL, White WB, et al.** Clinical practice guidelines for the management of hypertension in the community: a statement by the American Society of Hypertension and the International Society of Hypertension. *J Clin Hypertens (Greenwich)* 2014; 16:14–26.
17. **Ambrosius WT, Sink KM, Foy CG, et al; SPRINT Study Research Group.** The design and rationale of a multicenter clinical trial comparing two strategies for control of systolic blood pressure: the Systolic Blood Pressure Intervention Trial (SPRINT). *Clin Trials* 2014; 11:532–546.
18. **Myers MG, Godwin M, Dawes M, Kiss A, Tobe SW, Kaczorowski J.** Conventional versus automated measurement of blood pressure in the office (CAMBO) trial. *Fam Pract* 2012; 29:376–382.
19. **Messerli FH, Mancia G, Conti CR, et al.** Dogma disputed: can aggressively lowering blood pressure in hypertensive patients with coronary artery disease be dangerous? *Ann Intern Med* 2006; 144:884–893.
20. **Boutitie F, Gueyffier F, Pocock S, Fagard R, Boissel JP; INDANA Project Steering Committee; INdividual Data ANalysis of Antihypertensive intervention.** J-shaped relationship between blood pressure and mortality in hypertensive patients: new insights from a meta-analysis of individual-patient data. *Ann Intern Med* 2002; 136:438–448.
21. **Mancia G.** Effects of intensive blood pressure control in the management of patients with type 2 diabetes mellitus in the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial. *Circulation* 2010; 122:847–849.
22. **Perkovic V, Rodgers A.** Redefining blood-pressure targets—SPRINT starts the marathon. *N Engl J Med* 2015; 373:2175–2178.
23. **Margolis KL, O'Connor PJ, Morgan TM, et al.** Outcomes of combined cardiovascular risk factor management strategies in type 2 diabetes: the ACCORD randomized trial. *Diabetes Care* 2014; 37:1721–1728.
24. **Peterson JC, Adler S, Burkart JM, et al.** Blood pressure control, proteinuria, and the progression of renal disease. The Modification of Diet in Renal Disease study. *Ann Intern Med* 1995; 123:754–762.
25. **Agodoa LY, Appel L, Bakris GL, et al; African American Study of Kidney Disease and Hypertension (AASK) Study Group.** Effect of

ramipril vs amlodipine on renal outcomes in hypertensive nephrosclerosis: a randomized controlled trial. *JAMA* 2001; 285:2719–2728.

26. **Berl T, Hunsicker LG, Lewis JB, et al; Collaborative Study Group.** Impact of achieved blood pressure on cardiovascular outcomes in the Irbesartan Diabetic Nephropathy Trial. *J Am Soc Nephrol* 2005; 16:2170–2179.

27. **Pohl MA, Blumenthal S, Cordonnier DJ, et al.** Independent and additive impact of blood pressure control and angiotensin II receptor blockade on renal outcomes in the Irbesartan Diabetic Nephropathy

Trial: clinical implications and limitations. *J Am Soc Nephrol* 2005; 16:3027–3037.

28. **Bress AP, Tanner RM, Hess R, Colantonio LD, Shimbo D, Muntner P.** Generalizability of results from the Systolic Blood Pressure Intervention Trial (SPRINT) to the US adult population. *J Am Coll Cardiol* 2015 Oct 31. doi: 10.1016/j.jacc.2015.10.037. Epub ahead of print.

ADDRESS: George Thomas, MD, Department of Nephrology and Hypertension, Q7, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195; e-mail: thomasg3@ccf.org

CME CREDIT **CME CALENDAR**

2016

APRIL

CONTEMPORARY MANAGEMENT OF VALVULAR AND STRUCTURAL HEART DISEASE: PUTTING THE GUIDELINES AND TRIALS INTO PERSPECTIVE
April 1
Chicago, IL

AMBULATORY UROLOGY SYMPOSIUM
April 8
Cleveland, OH

WAKE UP TO SLEEP DISORDERS 2016: A CLEVELAND CLINIC SLEEP DISORDERS CENTER UPDATE
April 11–13
Cleveland, OH

MASTERS' APPROACH TO CRITICAL LIMB ISCHEMIA
April 17–19
Cleveland, OH

MAY

PRINCIPLES AND ADVANCES IN PERICARDIAL INTERVENTION IN CARDIAC ELECTROPHYSIOLOGY
May 3
San Francisco, CA

INTERNATIONAL LEAD MANAGEMENT COLLABORATIVE COMMUNITY SYMPOSIUM: A FRESH APPRAISAL

May 4
San Francisco, CA

21ST ANNUAL DIABETES DAY
May 6
Cleveland, OH

2016 FELLOWS' CONFERENCE: ADVANCES IN ORGAN TRANSPLANTATION
May 6–8
Cambridge, MA

21ST ANNUAL MAY DAY THERAPY: DERMATOLOGY
May 12
Cleveland, OH

SPASTICITY AND OTHER MOVEMENT DISORDERS: PEDIATRIC AND ADULT SYMPOSIUM
May 13–14
Warrensville Heights, OH

18TH ANNUAL INNOVATIONS IN NEUROSCIENCE
May 20
Cleveland, OH

ULTRASOUND: FROM BASICS TO BEDSIDE DIAGNOSIS AND PROCEDURES
May 20–21
Cleveland, OH

2016 INTERNATIONAL MOBILE STROKE SUMMIT: TRANSFORMING STROKE CARE THROUGH INNOVATIVE STROKE SYSTEMS

May 20–21
Cleveland, OH

JUNE

7TH ANNUAL INTERNAL MEDICINE BOARD REVIEW AND UPDATE
June 4–8
Sunrise, FL

28TH ANNUAL INTENSIVE REVIEW OF INTERNAL MEDICINE
June 5–10
Cleveland, OH

ABIM MAINTENANCE OF CERTIFICATION LEARNING SESSION
June 11
Cleveland, OH

BREAKTHROUGHS IN COGNITIVE NEUROLOGICAL DISORDERS
June 11
Las Vegas, NV

MELLEN CENTER UPDATE IN MULTIPLE SCLEROSIS
June 24
Cleveland, OH

AUGUST

PALLIATIVE MEDICINE AND SUPPORTIVE ONCOLOGY 2016
August 4–6
Cleveland, OH

FOR SCHEDULE UPDATES AND TO REGISTER, VISIT: WWW.CCFMEO.ORG